

Evidence of a Massive Black Hole Companion in the Massive Eclipsing Binary V Puppis

Qian S.-B.^{1,2,3}, Liao W.-P.^{1,2,3}, and Fernández Lajús, E.⁴

ABSTRACT

Up to now, most stellar-mass black holes were discovered in X-ray emitting binaries, in which the black holes are formed through a common-envelope evolution. Here we give evidence for the presence of a massive black hole candidate as a tertiary companion in the massive eclipsing binary V Puppis. We found that the orbital period of this short-period binary ($P=1.45$ days) shows a periodic variation while it undergoes a long-term increase. The cyclic period oscillation can be interpreted by the light-travel time effect via the presence of a third body with a mass no less than 10.4 solar mass. However, no spectral lines of the third body were discovered indicating that it is a massive black hole candidate. The black hole candidate may correspond to the weak X-ray source close to V Puppis discovered by Uhuru, Copernicus, and ROSAT satellites produced by accreting materials from the massive binary via stellar wind. The circumstellar matter with many heavy elements around this binary may be formed by the supernova explosion of the progenitor of the massive black hole. All of the observations suggest that a massive black hole is orbiting the massive close binary V Puppis with a period of 5.47 years. Meanwhile, we found the central close binary is undergoing slow mass transfer from the secondary to the primary star on a nuclear time scale of the secondary component, revealing that the system has passed through a rapid mass-transfer stage.

Subject headings: Stars: binaries : close – Stars: binaries : eclipsing — Stars: individuals (V Puppis) — Stars: evolution — Stars: black hole candidate

¹National Astronomical Observatories/Yunnan Observatory, Chinese Academy of Sciences (CAS), P.O. Box 110, 650011 Kunming, P.R. China (e-mail: qsb@netease.com)

²United Laboratory of Optical Astronomy, Chinese Academy of Sciences (ULOAC), 100012 Beijing, P. R. China

³Graduate School of the CAS, Beijing, P.R. China

⁴Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata, Paseo del Bosque s/n, 1900, La Plata, Pcia. Bs. As., Argentina

1. Introduction

With an orbital period of 1.4545 days and two B-type components stars (spectral types B1 and B3), V Puppis (HR 3129 and HD 65818) is one of a few massive semi-detached eclipsing binaries where the less massive components are filling their Roche lobes (e.g., Andersen et al. (1983); Bell et al. (1987a, b); Terrell et al. (2005)). According to the theory of binary evolution (e.g., Sybesma 1986), these systems were formed through a case A mass-transfer (mass transfer is occurring when hydrogen is burning in the core). Since the discovery of V Puppis more than one century ago, it was extensively investigated photometrically and spectroscopically (e.g. Andersen et al. (1983) and references therein). A weak X-ray source in the field of V Puppis (e.g., Giacconi et al. (1974); Groote et al. 1978); Bahcall et al. (1975)) and a significant amount of circumstellar matter around the binary (e.g., York et al. 1976 and Koch et al. 1981) make it a very interesting binary to study.

Because of its stable light curve and the deep and sharp eclipsing minima (e.g., Andersen et al. (1983)), times of light minimum of V Puppis can be determined precisely. Thus, the variations of the orbital period can in principle be derived with high precision by analyzing the observed-calculated (O-C) diagram. In the present letter, orbital period changes of V Puppis are investigated. Then, based on the period variations, the presence of a massive black hole companion and the evolutionary state of the system are discussed.

2. Analysis of the orbital period changes

Epochs and orbital periods of V Puppis have been given by several authors (e.g., Andersen et al. 1983). Kreiner & Ziolkowski (1978) did not detect any period change for the binary, while Andersen et al. (1983) pointed out that the period of V Puppis is variable. However, since the early eclipse times were mainly observed photographically or visually, conclusions on the period change are not reliable. For the present analysis, we only use the photoelectric (Pe) and charge-coupled device (CCD) observations. Most of the data were collected by Kreiner et al. (2001). Two new CCD times of light minimum were obtained on November 2, 2007 and on March 24, 2008 with the Virpi S. Niemela 0.8-m telescope at La Plata Observatory (see Table 1). During the observation, the V filter was used. The $(O - C)_1$ values of all available Pe and CCD times of light minimum were computed with the linear ephemeris given by Kreiner et al. (2001),

$$Min.I = JD(HEL.)2428648.2837 + 1^d.45448686 \times E, \quad (1)$$

where 2428648.2837 is the time of conjunction, $1^d.45448686$ the constant ephemeris period P_e , and E the cycle. The $(O - C)_1$ values are listed in the sixth column of Table 1 and are represented graphically against E in the upper panel of Figure 1.

If the orbital period of a binary varies linearly in time, the (O-C) diagram should have a parabolic shape, as observed in other eclipsing binaries (e.g., Qian & Zhu 2006). As for V Puppis, a simple second-order polynomial does not give a satisfactory fit (the dashed line in the upper panel of Figure 1). Therefore, a sinusoidal term was added to a quadratic ephemeris to give a good fit to the $(O - C)_1$ curve. A least-squares solution leads to the following ephemeris,

$$\begin{aligned} Min.I &= 2428648.2960(\pm 0.0045) + 1^d.45448114(\pm 0.00000091) \times E \\ &\quad + 4.18(\pm 0.44) \times 10^{-10} \times E^2 \\ &\quad + 0.0163(\pm 0.0016) \sin[0.^{\circ}2621(\pm 0.0011) \times E + 44.^{\circ}3(\pm 3.^{\circ}9)]. \end{aligned} \quad (2)$$

The sinusoidal term suggests an oscillation with a period of about $T = 5.47$ year and an amplitude of about $A = 0.^d0163$, which suggests a cyclic period oscillation with a period of 5.47 years (the upper panel of Fig. 1). Though the $(O - C)_1$ data do not cover a whole cycle, as shown in Figure 1, the minimum, the maximum, and the period of the cyclic variation are all well constrained, indicating that the period changes are reliable. The quadratic term in Eq. (2) reveals a continuous period increase at a rate of $dP/dt = +2.10 \times 10^{-7}$ days/year corresponding to a period increase of 1.8 s per century.

3. Discussion of the orbital period variation

3.1. The presence of a massive black hole companion

Both components of V Puppis are B-type stars containing a convective core and a radiative envelope, which is different from the Sun. Hence, the period oscillation can not be explained as solar-type magnetic activity cycles (i.e. the Applegate mechanism) (Applegate 1992). Therefore, the simplest explanation of the cyclic period oscillation is a wobble of the binary's barycentre due to the presence of a tertiary companion. The $(O - C)_2$ values calculated from the quadratic ephemeris in Eq. (2) are shown in the seventh column of Table 1. The corresponding $(O - C)_2$ diagram is plotted against the epoch number E in the middle panel of Figure 1. By considering a general case with an elliptical orbit, the following equation was used to describe those $(O - C)_2$ residuals,

$$(O - C)_2 = a_0 + \sum_{i=1}^2 [a_i \cos(i\Omega E) + b_i \sin(i\Omega E)]. \quad (3)$$

The results were obtained with the same value of Ω ($0.^\circ 2621$) as that used in equation (2) (solid line in the middle panel of Figure 1). The orbital parameters of the tertiary companion were computed by using the formulae derived by Kopal (1959),

$$a'_{12} \sin i' = c \sqrt{a_1^2 + b_1^2}, \quad (4)$$

$$e' = 2 \sqrt{\frac{a_2^2 + b_2^2}{a_1^2 + b_1^2}}, \quad (5)$$

$$\omega' = \arctan \frac{(b_1^2 - a_1^2)b_2 + 2a_1a_2b_1}{(a_1^2 - b_1^2)a_2 + 2a_1b_1b_2}, \quad (6)$$

$$\tau' = t_0 - \frac{T}{2\pi} \arctan \frac{a_1b_2 - b_1a_2}{a_1a_2 + b_1b_2}, \quad (7)$$

where c is the speed of light, a'_{12} , i' , e' , ω' , and τ' are the semi-major axis, the orbital inclination, the eccentricity, the longitude of the periastron from the ascending node, and the time of the periastron passage, respectively. The solutions are displayed in Table 2. With the following well-known equations,

$$f(m) = \frac{4\pi^2}{GT^2} \times (a_{12} \sin i')^3 \quad (8)$$

and

$$f(m) = \frac{(M_3 \sin i')^3}{(M_1 + M_2 + M_3)^2}, \quad (9)$$

the mass function, the masses and radii of the third body were calculated using the absolute parameters determined by Andersen et al. (1983). In the formula, M_1 , M_2 , and M_3 are the masses of the eclipsing pair and the third body, respectively, G is the gravitational constant, and T is the period of the (O-C) oscillation. For different values of the orbital inclination, the masses and orbital radii of the tertiary component are displayed in Table 2. The corresponding relations between the mass and the radius of the third body and the orbital inclination are shown in Figure 2.

The lowest mass of the tertiary companion is $10.4 M_\odot$, i.e. larger than the mass of the secondary component. If the tertiary component is a normal star, we would see its spectral lines not changing with the orbital phase of the binary. However, this has not been reported (e.g., York et al. 1976; Koch et al. 1981; Andersen et al. 1983). Moreover, the third body is more massive than the secondary component ($M_2 = 7.76 M_\odot$) in the central binary, it

should be very luminous and should contribute a large amount of third light to the total system unless it is not a normal main-sequence star. Therefore, we suspect that the tertiary component is a compact object and a probable black hole candidate. This situation resembles that in the triple system HR 2876 where a compact object (possibly a white dwarf star) with a mass excess of $1.0 M_{\odot}$ was found to be orbiting a B-type close binary star (B3.5V+B6V; P=15 days) (Burleigh & Barstow 1998; Vennes 2000). The main differences between the two systems are that the tertiary companion in the V Puppis is much more massive and that the central eclipsing binary in V Puppis system has a much more tight orbit and shows strong interaction between both components.

A weak X-ray source (3U 0750-49) with 9.4 ± 2.3 count s^{-1} was detected by the Uhuru satellite (Giacconi et al. 1974). The counterpart of the X-ray source was a puzzling problem for astronomers since the 70s of last century. V Puppis was initially listed in the 3U catalogue as a possible candidate by Giacconi et al. (1974). However, another bright star HD 64740, close to the source (within the 90 percent confidence error box) was also considered as a possible candidate (Groote et al. 1978). By analyzing the X-ray data of V Puppis and HD 64740 from the Copernicus satellite, Bahcall et al. (1975) later concluded that the weak X-ray source must almost certainly be closer to V Puppis than to HD 64740, but most probably neither V Puppis nor HD 64740 is the candidate. More recently, V Puppis was also discovered to be a X-ray source by ROSAT satellite with 0.08 count s^{-1} (X-ray 50 Å, 250 eV) (e.g., Lampton et al. 1996). Our study indicates that the tertiary companion may correspond to the weak X-ray source. There are two mechanisms that can produce the X-ray emission in the V Puppis system. If the massive tertiary companion is a normal star, the X-ray emission could be caused by the colliding winds from the eclipsing pair and the tertiary companion, resembling those observed in other massive binary systems (e.g., Stevens 1992; Pittard & Stevens 2002). If the third body is a massive black hole, the X-ray source is reproduced by accreting material from the massive binary via stellar wind as observed in X-ray emitting binaries (e.g., Prestwich et al. 2007). However, the former mechanism can be ruled out, because no spectral lines of the tertiary component were found.

Another interesting feature of V Puppis is the existence of a significant amount of circumstellar matter, which was deduced from the optical spectra and ultraviolet observations obtained by the IUE spacecraft and the Copernicus satellite (e.g., York et al. 1976; Koch et al. 1981; Andersen et al. 1983). Numerous interstellar spectral lines, especially those of many types of heavy elements (e.g., Mg II, Si I, Si II, Si III, C I, O I, Fe II and Fe III) were found in the H II region around V Puppis, in which strong lines of Si III and Fe III are remarkable (e.g., York et al. 1976 and Koch et al. 1981). The formation of the circumstellar matter around V Puppis is an unsolved problem. Considering that V Puppis has undergone mass exchange, one may think that the circumstellar matter results from from the binary

evolution. However, we may question why the other systems (e.g. AI Cru) that resemble V Puppis do not have the same significant amount of circumstellar matter (Bell et al. 1987). The most probable reason for the formation of the circumstellar matter around V Puppis is that it is formed by the supernova explosion of the progenitor of the tertiary black hole companion (e.g., Woosley & Weaver (1995) and Zhang et al. (2007)). This may produce the heavy elements in the circumstellar matter and results in the formation of the H II region around the eclipsing binary.

3.2. The slow mass transfer of V Puppis

V Puppis is a semi-detached binary system with the secondary component filling the critical Roche lobe (e.g., Andersen et al. 1983). The long-term period increase can be explained by the mass transfer from the secondary to the primary component. By using the equation,

$$\tau_N = 10^{10} M_2 / L_2, \quad (10)$$

where M_2 and L_2 are the masses and the luminosity of the less massive component, the nuclear time-scale of the secondary is calculated to be $\tau_N = 7.1 \times 10^6$ years. The value is close to the timescale of the period change $\tau_P = \frac{P}{\dot{P}} = 6.9 \times 10^6$ years, which reveals that V Puppis is now undergoing a slow mass-transfer evolutionary stage on the nuclear time-scale of the less massive component. This suggests that V Puppis is formed via a case A evolution and has passed through a rapid mass-transfer stage. The results are in agreement with the prediction from the theory of massive binary evolution (e.g., Sybesma 1986). By assuming a conservative mass transfer, the well-known equation

$$\frac{\dot{P}}{P} = 3\dot{M}\left(\frac{1}{M_2} - \frac{1}{M_1}\right), \quad (11)$$

yields a rate of mass transfer $dM/dt = 7.82 \times 10^{-7} M_\odot/\text{year}$. However, since the nuclear time-scale of the secondary is slightly longer than the time scale of period change, the slow mass transfer is insufficient to cause the observed period increase. This suggests that the stellar wind from the massive component of the binary should contribute to the period increase. The situation resembles that of AI Cru (Qian et al. 2008).

By considering conservative mass and angular momentum, the calculation by Plavec (1968) suggests that, to produce a case A mass transfer of massive binaries, the initial orbital period should be less than 1.8 days. One may ask a question what is the reason that causes the origin of the initially short-period detached system? It is possible that the massive tertiary companion has played an important role for the origin and evolution of inner binary

by removing angular momentum from the central system via Kozai oscillation (Kozai 1962) or a combination of Kozai cycle and tidal friction (e.g., Fabrycky & Tremaine 2007). This makes the inner eclipsing pair to have a lower angular momentum and a shorter initial orbital period.

4. Conclusions

The cyclic change of the orbital period, the presence of a weak X-ray source near V Puppis, and the significant amount of circumstellar matter around the binary star all support the conclusion that a massive black hole is orbiting the massive close binary V Puppis with a period of 5.47 years. Several massive stellar black hole candidates have been discovered recently (e.g., Greiner et al. (2001) and Orosz (2007)), but all candidates are found to be a component in a binary system. We conclude here that we have provided observational evidence for the first massive stellar black hole candidate as a tertiary companion of a massive close binary star. Theoretical investigations have shown that the formation of black hole binaries should experience a common-envelope phase with the spiral-in of the companion in the envelope causing the ejection of the envelope (e.g., Taam & Sandquist (2000) and Podsiadlowski et al. (2002)). The mean distance between the black hole candidate and the eclipsing pair is about 5.5 astronomical units (AU). During the supergiant phase, the radius of the precursor of the black hole companion in V Puppis may reach to several AUs. The survival of V Puppis in this evolutionary stage suggests that the tertiary black hole companion must be formed through an evolutionary process that is different from that of the black holes in X-ray binaries (e.g., Woosley & Weaver (1995) and Zhang et al. (2007)) and it could give some constraints to the evolution of single massive stars and the formation of isolated black holes.

From the present investigation, we deduce that the progenitor of V Puppis is a triple system that is composed of a short-period detached B-type eclipsing binary and a much more massive tertiary companion. By drawing angular momentum from the inner binary through Kozai oscillation (Kozai 1962) or a combination of Kozai cycle and tidal friction (e.g., Fabrycky & Tremaine 2007), the massive tertiary companion may play an important role for the origin the central system, which causes the inner eclipsing binary to have a very short initial orbital period. In the triple system, the massive tertiary companion evolves faster, and finally into a massive black hole via a supernova explosion (e.g., Woosley & Weaver (1995) and Zhang et al. (2008)). At the same time, the original more massive component of the eclipsing pair evolves to fill its critical Roche lobe. Then, the binary was undergoing a rapid mass-transfer stage, and after the mass ratio was reversed, the central binary finally reached

the present semi-detached configuration with a slow secondary-to-primary mass transfer on the nuclear time scale of the present less massive star. Supergiant stars have recently been found to be tertiary companions of OB-type close binaries in the two triple systems HD 167971 and FR Sct (e.g., Leitherer et al. (1987); Davidge & Forbes (1988), and Pigulski & Michalska (2007)). They resemble the progenitor of V Puppis with the tertiary supergiant companion evolving into a compact object. As for V Puppis, the existence of the black hole in the outer orbit poses strong constrain on the formation of the black hole, the mass loss in the supernova that lead to the formation of the black hole, and the magnitude of the kick received by the black hole, all make the object a much more interesting system for future study.

This work is partly supported by the Yunnan Natural Science Foundation (No. 2005A0059M), Chinese Natural Science Foundation (No.10573032 and No.10433030), and the National Key Fundamental Research Project through grant 2007CB815406. New CCD photometric observations of V Puppis were obtained with the Virpi S. Niemela 0.8-m telescope at La Plata Observatory. We are grateful to Prof. Kreiner for sending us the times of light minimum of the binary and to Cecilia Fariña and Carolina von Essen for their helps during the observations and the data processing. The authors thank Prof. J.-P. De Greve for improving the English writing of the original manuscript and the referee for those useful comments and suggestions that help to improve the paper.

REFERENCES

- Andersen, J., et al., 1983, A&A 128, 17
- Applegate, J. H. A, 1992, ApJ 385, 621
- Bahcall, J. N., et al., 1975, MNRAS 171, 41
- Bell, S. A., Adamson, A. J., Hilditch, R. W., 1987a, MNRAS 224, 649
- Bell, S. A., Kilkenny, D., Malcolm, G. J., 1987b, MNRAS 226, 879
- Burleigh, M. R. & Barstow, M. A., 1998, MNRAS 295, L15
- Charles, P. A. & Coe, M. J., 2006, In: Compact stellar X-ray sources (Eds. by W. Lewin & M. van der Klis), P. 215 (Cambridge University Press, Cambridge)
- Davidge, T. J. & Forbes, D., 1988, MNRAS 235, 797
- Fabrycky, D. & Tremaine, S., 2007, ApJ 669, 1298
- Giacconi, R. et al., 1974, ApJS 27, 37

- Greiner, J., Cuby, J. G., McCaughrean, M. J., 2001, *Nature* 414, 522
- Groote, D., Kaufmann, J. P., 1978, *A&A* 63, L9
- Koch, R. H. et al., 1981, *PASP* 93, 621
- Kopal, Z., 1959, *Close Binary systems*. (London: Chapman & Hall). P.112
- Kozai, Y., 1962, *AJ* 67, 591
- Kreiner, M. J., 2006, private communication.
- Kreiner, M. J., Kim, C.-H., Nha, I.-S, 2001, *An Atlas of O-C Diagrams of Eclipsing Binaries*, Krakov, P.1914-1915
- Kreiner, M. J. & Ziolkowski, J., 1978, *Acta Astron.* 28, 497
- Lampton, M., Lieu, R., Schmitt, J. H. M. M., Bowyer, S., Voges, W., Lewis, J., and Wu, X., 1997, *ApJS* 108, 545
- Leitherer, C. et al., 1987, *A&A* 185, 121
- Orosz, J. A. A, 2007, *Nature* 449, 872
- Pigulski, A. & Michalska, G., 2007, *Inf. Bull. Var. Stars* No. 5757
- Pittard, J. M. & Stevens, I. R., 2002, *A&A* 388, L20
- Plavec, M., 1968, *Adv. Astr. Astrophys.* 6, 202
- Podsiadlowski, Ph., Rappaport, S., Han, Z., 2002, *MNRAS* 341, 385
- Prestwich, A. H., et al., 2007, *ApJ* 669, L21
- Qian S.-B., Zhao, E.-G., Fernández Lajús, E., Zhu L.-Y., Liu L., Liao W.-P., Carolina von E., 2008, in preparation
- Qian, S.-B. & Zhu, L.-Y., 2006, *AJ* 131, 1032
- Remillard, R. A. & McClintock, J. E., 2006, *Annu. Rev. Astron. Astrophys.* 44, 49
- Stevens, I. R., Blondin, J. M., Pollock, A. M. T., 1992, *ApJ* 386, 265
- Stickland, D. J., Lloyd, C., Pachoulakis, I., Koch, R. H., 1998, *Observatory* 118, 356
- Sybesma, C. H. B., 1986, *A&A* 159, 108
- Taam, R. E. & Sandquist, E. L., 2000, *Annu. Rev. Astron. Astrophys.* 38, 113
- Terrell, D., Munari, U., Zwitter, T., Wolf, G., 2005, *MNRAS* 360, 583
- Vennes, S., 2000, *A&A* 354, 995
- Woosley, S. E. & Weaver, T. A., 1995, *ApJS* 101, 181
- York, D. G., Flannery, B., Bahcall, J., 1976, *ApJ* 210, 143

Zhang, W.-Q., Woosley, S. E., Heger, A., 2008, ApJ 679, 639

Table 1: All available photoelectric and CCD times of light minimum for V Puppis.

JD.Hel. 2400000+	Errors days	Meth.	Min.	E	$(O - C)_1$ days	$(O - C)_2$ days	Residuals days	Ref.
31917.957	± 0.0010	Pe	I	2248	-0.0132	-0.0147	-0.0004	(1)
31965.952	± 0.0010	Pe	I	2281	-0.0162	-0.0176	-0.0046	(1)
39869.6264	± 0.0010	Pe	I	7715	-0.0234	-0.0164	-0.0011	(1)
41791.0113	± 0.0002	Pe	I	9036	-0.0157	-0.0104	+0.0065	(2)
42537.9043	± 0.0001	Pe	II	9549.5	-0.0017	+0.0026	-0.0042	(2)
42561.9044	± 0.0005	Pe	I	9566	-0.0006	+0.0036	-0.0044	(2)
43124.8032	± 0.0002	Pe	I	9953	+0.0118	+0.0151	-0.0001	(2)
43148.8024	± 0.0002	Pe	II	9969.5	+0.0119	+0.0151	+0.0014	(2)
43151.7111	± 0.0002	Pe	II	9971.5	+0.0117	+0.0149	+0.0014	(2)
45367.6063	± 0.0002	Pe	I	11495	-0.0039	-0.0056	-0.0022	(2)
45383.6052	± 0.0001	Pe	I	11506	-0.0043	-0.0061	-0.0014	(2)
48391.5025	± 0.0010	CCD	I	13574	+0.0142	+0.0026	+0.0021	(3)
48425.6868	± 0.0007	CCD	II	13597.5	+0.0180	+0.0063	+0.0044	(3)
48500.5947	± 0.0050	Pe	I	13649	+0.0198	+0.0078	+0.0025	(4)
49821.2500	± 0.0050	Pe	I	14557	+0.0011	-0.0164	-0.0003	(4)
54456.7420	± 0.0010	CCD	I	17744	+0.0435	+0.0012	-0.0024	(5)
54550.5667	± 0.0010	CCD	II	17808.5	+0.0538	+0.0109	+0.0028	(5)

References in Table 1:

(1) Kreiner & Ziolkowski (1978); (2) Andersen et al. (1983); (3) Kreiner (2006); (4) Stickland (1998); (5) The present authors.

Table 2: Calculated orbital parameters of the tertiary companion in V Puppis.

Parameters	Values	Errors
a_0	-0.0005	± 0.0001
a_1	+0.0142	± 0.0012
b_1	+0.0113	± 0.0012
a_2	-0.0031	± 0.0019
b_2	+0.0026	± 0.0004
$a'_{12} \sin i'$ (AU)	3.14	± 0.21
e'	0.45	± 0.16
ω'	$-63.^{\circ}0$	$\pm 18.^{\circ}9$
τ' (days)	2428213	± 101
$f(m)$ ($M_{\odot}$)	1.03	± 0.21
$M_3(i' = 90^{\circ})$	10.40	± 0.87
$M_3(i' = 70^{\circ})$	11.25	± 0.96
$M_3(i' = 50^{\circ})$	14.74	± 1.33
$M_3(i' = 30^{\circ})$	27.44	± 2.87
$d_3(i' = 90^{\circ})$	6.82	± 0.73
$d_3(i' = 70^{\circ})$	6.80	± 0.73
$d_3(i' = 50^{\circ})$	6.28	± 0.70
$d_3(i' = 30^{\circ})$	5.56	± 0.66




Fig. 1.— A plot of the $(O - C)_1$ curve of V Puppis with respect to the linear ephemeris given by Kreiner et al. (2001). The solid line in the upper panel suggests a combination of a long-term period increase and a cyclic change, while the dashed line refers to the continuous increase of the orbital period. The middle panel shows the plot of the $(O - C)_2$ residuals of the eclipse timings based on the quadratic ephemeris in Eq. (1). The solid line in the middle panel is the theoretical orbit of the massive black hole candidate with an eccentricity of 0.45. Residuals after subtracting all effects of period change are displayed in the lower panel, and no variations can be traced there.




Fig. 2.— The relations between the mass $M_3 (M_\odot)$ and the orbital radius d_3 (AU) of the tertiary component and its orbital inclination i' in the V Puppis system. The tertiary companion should be more massive than the secondary component and thus may be a black hole candidate.